

POSTER PRESENTATION

Open Access

# Continuous or discrete? Attractor dynamics and spatial representations in a model of the hippocampal network

Federico Stella<sup>1\*</sup>, Remi Monasson<sup>2</sup>, Alessandro Treves<sup>1,3</sup>

From Twenty Second Annual Computational Neuroscience Meeting: CNS\*2013  
Paris, France. 13-18 July 2013

There is now strong evidence of the presence of a vast integrated neural system, which involves the hippocampus and several parahippocampal regions, dedicated to spatial navigation and memory. 'Place cells' and 'Grid cells' have been identified as neural correlates of the ability to remember an environment and to successfully move through it.

The storage of a spatial map in the hippocampal circuits poses important computational issues. It asks for an extension of the classical notion of attractor, successfully used to model the storage of discrete memory items, like episodic memories, to a new entity that captures the continuous nature of space.

In a continuous attractor activity smoothly changes with the position of the animal in the environment, so that each point in space is uniquely represented in the hippocampus. This situation corresponds to the presence of an infinite number of distinct stable configurations of the system, each for a different point of the environment.

It is not known how much this idealized notion corresponds to the actual structure of the maps stored in the hippocampus, and how well attractor dynamics can approximate a continuous representation.

We address these questions, within a simplified mathematical network model. The model network simulates the storage in CA3 of a spatial representation, its retrieval and its transfer to CA1. Through both analytical calculations and computer simulations we evaluate the ability of the network to represent different points in space and we study the properties of the attractor landscape when different kinds of disorder (connections

dilution, heterogeneous place field distribution, multiple maps) are introduced in the system. Moreover we investigate the possible role of CA1 in the realization of a continuous attractor.

We find that even networks of considerable size can only approximate the idealized notion of a 2D quasi-continuous dynamical attractor, but that the presence of CA1 generates a 'smoother' representation of space.

#### Author details

<sup>1</sup>Cognitive Neuroscience Sector, Sissa, Trieste, Italy. <sup>2</sup>Laboratoire de Physique Theorique, ENS, Paris, France. <sup>3</sup>Embassy of Italy, Science Office, Tel Aviv, Israel.

Published: 8 July 2013

doi:10.1186/1471-2202-14-S1-P397

**Cite this article as:** Stella et al.: Continuous or discrete? Attractor dynamics and spatial representations in a model of the hippocampal network. *BMC Neuroscience* 2013 **14**(Suppl 1):P397.

#### Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Cognitive Neuroscience Sector, Sissa, Trieste, Italy  
Full list of author information is available at the end of the article