

Poster presentation

Open Access

Irregular persistent activity induced by synaptic excitatory feedback

Francesca Barbieri*¹ and Nicolas Brunel^{1,2}

Address: ¹Computational Neuroscience Unit, I.S.I. Foundation, Torino, Italy and ²Laboratory of Neurophysics and Physiology, UMR 8119 CNRS-Université René Descartes, Paris, France

Email: Francesca Barbieri* - barbieri@isiosf.isi.it

* Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, **8**(Suppl 2):P167 doi:10.1186/1471-2202-8-S2-P167

© 2007 Barbieri and Brunel; licensee BioMed Central Ltd.

Neuro-physiological experiments on monkeys [1] have reported highly irregular persistent activity during the performance of an oculomotor delayed-response task. These experiments show that during the delay period the ISI's coefficient of variation (CV) of prefrontal neurons is above 1, on average, and larger than during the fixation period, regardless of whether the cue is preferred or non-preferred.

Previous models [2,3] of spontaneous and selective persistent activity in the cortex based on excitatory synaptic feedback do not reproduce this feature because the excitatory feedback during persistent activity brings neurons in a region of the f-I curve in which the firing is relatively independent from fluctuations and hence the CV is small. To overcome this problem, we introduced two ingredients: (1) a high post-spike reset potential (close to threshold), (2) a non-linear relationship between synaptic efficacy and pre-synaptic firing rate via a short-term depression (STD) mechanism.

We show that when the reset potential is close enough to the threshold, the CV-I curve has a maximum above 1 for a sub-threshold mean current. The range of the mean synaptic input values for which the CV is greater than 1 is always in the sub-threshold regime in which firing is dominated by fluctuations of the mean synaptic input. With short-term depression, synaptic efficacies saturate at a certain limiting value of the presynaptic frequency; this in turn provokes a saturation of the mean synaptic current to a neuron at the same limiting presynaptic frequency. This

allows the persistent state solution to reach the region of the f-I curve which corresponds to high values of the CV.

We tested this idea both with numerical simulations and analytical techniques. For the analytical studies we used mean-field techniques, recently extended in presence of STD [4], that involves the use of the distribution of the interspike intervals of an integrate-and-fire neuron receiving a Gaussian current in input; this permits to obtain an accurate estimate of the statistic of the postsynaptic current in presence of STD and hence to find the stationary states in a self-consistent way. We also simulated both a fully connected excitatory network of leaky integrate-and-fire neurons endowed with STD, and a cortical network model composed of an inhibitory population and several stimulus selective excitatory populations.

In both cases we find a large range of values of the synaptic efficacies for which the persistent activity is irregular, with values of the CV in agreement with the physiological findings.

References

1. Compte A, Constantinidis JT, Raghavachari S, Chafee MV, Goldman-Rakic PS, Wang XJ: **Temporally irregular mnemonic persistent activity in prefrontal neurons of monkeys during a delayed response task.** *J Neurophysiol* 2003, **90**:3441-3454.
2. Amit DJ, Brunel N: **Model of global spontaneous activity and local structured activity during delay periods in the cerebral cortex.** *Cerebral Cortex* 1997, **7**:237-252.
3. Brunel N: **Persistent activity and the single-cell frequency-current curve in a cortical network model.** *Comput Neural Syst* 2000, **11**:1-20.

- Romani S, Amit DJ, Mongillo G: **Mean-field analysis of selective persistent activity in presence of short-term synaptic depression.** *J Comput Neurosci* **20**:201-217.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

