

Poster presentation

Open Access

Modeling perceptual multi-stability with Hodgkin-Huxley neurons

David Chik* and Roman Borisyuk

Address: Centre for Theoretical and Computational Neuroscience, University of Plymouth, Plymouth

Email: David Chik* - david.chik@plymouth.ac.uk

* Corresponding author

from Seventeenth Annual Computational Neuroscience Meeting: CNS*2008
Portland, OR, USA. 19–24 July 2008

Published: 11 July 2008

BMC Neuroscience 2008, 9(Suppl 1):P152 doi:10.1186/1471-2202-9-S1-P152

This abstract is available from: <http://www.biomedcentral.com/1471-2202/9/S1/P152>

© 2008 Chik and Borisyuk; licensee BioMed Central Ltd.

Perceptual multi-stability can be induced by either using some ambiguous figures (such as Necker Cube or Face/Vase figure) or showing different images to left and right eyes (binocular rivalry). The observer will experience a persistent switching between different perceptual representations of the same image. The switching process is spontaneous and irregular: time intervals between consequent switching would form some Gaussian-like distribution [1,2]. Switching is highly involuntary and usually human subjects cannot use their "will power" to control the process. Many behavioral experiments have been carried out. Levelt [3,4] found that during binocular rivalry, when the image brightness to one eye increased, the mean duration of perception from the other eye decreased (but little was change on the duration from the same eye). Leopold et al. [5] found that for ambiguous figures, the switching process would slow down if the figure appeared and disappeared periodically.

The neural mechanism of perceptual multi-stability is still unclear. It is possible that perceptual multi-stability is regulated in a similar way like selection process in object-based attention [6]. A very important discovery has been found from EEG recordings: the perception of ambiguous figures seems to correlate with gamma band activity and transient synchrony of some distant brain areas [7-10]. The idea of partial synchronization between a central element and some peripheral elements has been proposed for modeling selective attention [11]. This idea is supported by results of EEG recordings, and there is a strong suggestion that multi-stable perception might also be controlled by the mechanism of partial synchronization.

In the present study, we propose a model of cortical neural network for studying perceptual multi-stability. The model consists of central and peripheral neurons, all of which are modeled by Hodgkin-Huxley equations with synaptic couplings. A dynamical regime is found where initially the central neurons form partial synchronization with one group of peripheral neurons, but after some time, it spontaneously switches to synchronize with another group of peripheral neurons, and then switch back and forth. The model produces a similar Gaussian-like histogram as experimental data. It also meets the constraints observed from various experiments [3-5]. The synchronization occurs in the gamma range, in agreement with EEG findings [7-10].

References

1. Fox R, Herrmann J: **Stochastic properties of binocular rivalry alternations.** *Percept Psychophys* 1967, **2**:432-436.
2. Borsellino A, de Marco A, Allazetta A, Rinesi S, Bartolini B: **Reversal time distribution in the perception of visual ambiguous stimuli.** *Biological Cybernetics* 1972, **10**:139-144.
3. Levelt WJM: *On binocular rivalry. Minor Series 2. Psychological Studies The Hague: Mouton*; 1968.
4. Bossink CJH, Stalmeier PFM, de Weert CMM: **A test of Levelt's second proposition for binocular rivalry.** *Vision Res* 1993, **33**:1413-1419.
5. Leopold DA, Wilke M, Maier A, Logothetis NK: **Stable perception of visually ambiguous patterns.** *Nature Neurosci* 2002, **5**(6):605-609.
6. Mitchell JF, Stoner GR, Reynolds JH: **Object-based attention determines dominance in binocular rivalry.** *Nature* 2004, **429**:410-413.
7. Keil A, Muller MM, Ray WJ, Gruber T, Elbert T: **Human gamma band activity and perception of a Gestalt.** *J Neurosci* 1999, **19**(16):7152-7161.
8. Klemm WR, Li TH, Hernandez: **Coherent EEG indicators of cognitive binding during ambiguous figure tasks.** *Consciousness and Cognition* 2000, **9**:66-85.

9. Nakatani H, van Leeuwen C: **Transient synchrony of distant brain areas and perceptual switching in ambiguous figures.** *Biol Cybern* 2006, **94**:445-457.
10. Melloni L, Molina C, Pena M, Torres D, Singer W, Rodriguez E: **Synchronization of neural activity across cortical areas correlates with conscious perception.** *J Neurosci* 2007, **27(11)**:2858-2865.
11. Borisyuk RM, Kazanovich YB: **Oscillatory neural network model of attention focus formation and control.** *BioSystems* 2003, **71**:29-38.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

